

Match 6 Round 1 Geometry: Lines and Angles

Note: Figures not necessarily Drawn to scale




1.) What is the degree measure of the acute angle formed by the intersection of the lines { $y = \sqrt{3}(x - 2) + 1$, y - 4 = x - 5, x + y = 3} and $3y - 3 = \sqrt{3}(x - 2)$?

2.) In the figure below, segment AB is parallel to segment CD. The lines are cut by transversal line EF, which intersects line segment AB at G and segment CD at H. There is a number x such that the measure of angle AGE is $\{(\frac{5}{2}x - 23), (\frac{5}{2}x - 42), (\frac{5}{2}x - 61)\}$ degrees and the measure of angle GHD is $(\frac{2}{3}x + 89)$ degrees. Find the measure of angle FHD.



3.) Rhombus WXYZ has W at {(-4,3), (-8,6), (-6,8) } and X at (0,0). Y is in quadrant I and the slope of line XY is $\{\frac{1}{2}, \frac{1}{3}, \frac{1}{3}\}$. The slope of XZ is $\{a + b\sqrt{5}, a + b\sqrt{10}, a + b\sqrt{10}\}$ for some integer values of a and b. Find a+b.



1.) If the equation $\frac{1}{2}(x+3y-8) - z = 3z - 2(4x-y)$ is solved for z in terms of x and y, then z=ax+by+c. Find $\{a+b+c, a+b-c, a+b-2c\}$.

2.) If the equation { $x = \frac{y+1}{2}$, $x = y + \frac{1}{y}$, $x = \frac{y+1}{4}$ is solved for y, $y = \frac{ax \pm \sqrt{bx^2 + c}}{2}$ for some values of a, b, and c. Find a + b - c.

3.) Suppose that x > 0. When the equation $\begin{cases} 2xy(x^2 + 1) - 3x^2 = (2x^2 + 1)(2x^2 - 1) - x^2y - y, & 4xy(x^2 + 1) - 15x^2 = (4x^2 + 1)(4x^2 - 1) - x^2y - y, & 3xy(x^2 + 1) - 8x^2 = (3x^2 + 1)(3x^2 - 1) - x^2y - y \end{cases}$ is solved for y, the result is y = ax + b, for some constants a and b. Find a + b.

Match 6 Round 3 Geometry: Solids and Volumes

1.)_____

2.)_____

3.) _____

1.) A cone has horizontal base and its vertex lies vertically above the center of the base. The cone has height {8, 4, 12}, and its volume is {96 π , 12 π , 324 π }. The lateral area of the cone (the surface area not including the base) is A π . What is A?

2.) A sphere of radius {9, 6, 3} cm is inscribed in a cube. The volume that is outside the sphere but inside the cube is $a - b\pi$ cm³. What is a - b?

3.) A pyramid has a square base, and the base is horizontal. The height of the pyramid is {4, 8, 6}. A horizontal plane cuts the pyramid into two parts such that the volume of the top part is $\frac{1}{2}$ of the volume of the bottom part. If the height of this plane above the base is k, then $k = a - b\sqrt[3]{c}$, where a and b are rational numbers and c is an integer that is not divisible by the cube of any prime number. Find the product of a, b, and c.

Match 6 Round 4 Radical Expressions and Equations

1.	

2._____

3._____

1.)_ For how many integer values of *K* is $\{2 + \sqrt{K+3} - K, 4 + \sqrt{K+3} - K, 6 + \sqrt{K+3} - K\}$ a positive number?

2.) Suppose that a_0 , a_1 , a_2 ,... is a sequence of numbers such that $a_0 = x$, $a_1 = \sqrt{a_0}$, $a_2 = \sqrt{a_1}$, $a_3 = \sqrt{a_2}$, and so on. If $a_6 = \{2^7, 2^6, 2^9\}$, then $x = 2^n$. Find n.

3.) For how many integers *n* with $0 \le n \le \{10000,9000,8000\}$ is $\sqrt{2n+1}$ rational?

Match 6 Round 5 Polynomials									
and Advanced Factoring									
1									

- 2. _____
- 3.
- 1.) Let $f(x) = x^3 + Ax + B$ and suppose that $f(1) = \{3,5,7\}$ and $f(2) = \{15,20,25\}$. Find |A| + |B|.

2.

$$\{x^4 - 5x^3 + 17x^2 - 45x + K, x^4 - 5x^3 + 24x^2 - 80x + K, x^4 - 5x^3 + 12x^2 - 20x + K\}$$

factors to

 $\{(x^2 + 9)(x^2 + Bx + C), (x^2 + 16)(x^2 + Bx + C), (x^2 + 4)(x^2 + Bx + C)\}.$ Find K+B+C.

3.

A quartic polynomial $x^4 + Ax^3 + Bx^2 + Cx + D$, where *A*, *B*, *C*, *D* are integers, has {2+i and 1-2i , 3+i and 1-3i, 4+i and 1-4i} as two of its zeros, where $i = \sqrt{-1}$. Find A+B+C+D.

Match 6 Round 6 Counting and Probability


1.) _	 	 	
2.) _		 	
3.)			

1.) ${}_{N}C_{R}$ denotes the number of combinations of *N* objects taken *R* at a time. For how many of the {10,13,15} integer values of *R*, $0 \le R \le$ {9,12,14}, is { ${}_{9}C_{R}, {}_{12}C_{R,14}C_{R}$ } divisible by {9,12,14} ?

2.) The $\{12,10,14\}$ members of a club consist of $\{6,5,7\}$ married couples. A subset of 4 club members will be selected to represent the club at a conference. In how many ways can this be done if no person and his/her spouse may both be selected?

3) {Four nickels and six dimes, Three nickels and six dimes, Four nickels and five dimes} are placed in a bag, and five coins are drawn from the bag at random without replacement. The probability that the value of the coins is at least 40 cents is $\{\frac{A}{42}, \frac{A}{42}, \frac{A}{14}\}$. Find A.

Match 6 Team Round



1.) A, B, C, and D are the interior angles of a convex quadrilateral ABCD. The measure of the supplement of angle D is six degrees more than the measure of angle B. Find the sum of angles A and C.

2.) If
$$k = \sqrt[3]{3 + \sqrt[3]{3 + \sqrt[3]{3 + \sqrt[3]{3 + \dots}}}}$$
 and k is real, what is 100k rounded to the

nearest integer?

3.) A regular tetrahedron has volume $\frac{16\sqrt{2}}{3}$. The surface area is $K\sqrt{3}$ for some value of K. Find K.

4.) $x^3 + Ax^2 + Bx - 6$ factors into three binomials with integer coefficients. What is the absolute value of the sum of all possible values of A?

5) When three standard six-sided dice with sides labeled 1 through 6 are rolled, the probability that the sum is 12 or 13 is $\frac{K}{108}$. Find *K*.

6) A softball player has a probability 0.4 of getting a hit in any at bat. She comes to bat 5 times in one game, and the results of her at-bats are independent. The probability that she gets at least 3 hits is K. Find 100,000*K.